

1993). Such transcription units can be incorporated into a variety of vectors for introduction into mammalian cells, including but not restricted to, plasmid DNA vectors, viral DNA vectors (such as adenovirus or adeno-associated vectors), or viral RNA vectors (such as retroviral, semliki forest virus, sindbis virus vectors).

5 Ribozymes of this invention may be used as diagnostic tools to examine genetic drift and mutations within cell lines or cell types. They can also be used to assess levels of the target RNA molecule. The close relationship between ribozyme activity and the structure of the target RNA allows the detection of mutations in any region of the molecule which alters the base-pairing and three-dimensional structure of the target RNA.
10 By using multiple ribozymes described in this invention, one may map nucleotide changes which are important to RNA structure and function *in vitro*, as well as in cells and tissues. Cleavage of target RNAs with ribozymes may be used to inhibit gene expression and define the role (essentially) of specified gene products in particular cells or cell types.

15 4.14 ISOLATING HOMOLOGOUS GENE AND GENE FRAGMENTS

 The genes and δ -endotoxins according to the subject invention include not only the full-length sequences disclosed herein but also fragments of these sequences, or fusion proteins, which retain the characteristic insecticidal activity of the sequences specifically exemplified herein.
20

 It should be apparent to a person skill in this art that insecticidal δ -endotoxins can be identified and obtained through several means. The specific genes, or portions thereof, may be obtained from a culture depository, or constructed synthetically, for example, by use of a gene machine. Variations of these genes may be readily constructed using
25 standard techniques for making point mutations. Also, fragments of these genes can be made using commercially available exonucleases or endonucleases according to standard procedures. For example, enzymes such as *Bal31* or site-directed mutagenesis can be used to systematically cut off nucleotides from the ends of these genes. Also, genes which code for active fragments may be obtained using a variety of other restriction

enzymes. Proteases may be used to directly obtain active fragments of these δ -endotoxins.

Equivalent δ -endotoxins and/or genes encoding these equivalent δ -endotoxins can also be isolated from *Bacillus* strains and/or DNA libraries using the teachings provided herein. For example, antibodies to the δ -endotoxins disclosed and claimed herein can be used to identify and isolate other δ -endotoxins from a mixture of proteins. Specifically, antibodies may be raised to the portions of the δ -endotoxins which are most constant and most distinct from other *B. thuringiensis* δ -endotoxins. These antibodies can then be used to specifically identify equivalent δ -endotoxins with the characteristic insecticidal activity by immunoprecipitation, enzyme linked immunoassay (ELISA), or Western blotting.

A further method for identifying the δ -endotoxins and genes of the subject invention is through the use of oligonucleotide probes. These probes are nucleotide sequences having a detectable label. As is well known in the art, if the probe molecule and nucleic acid sample hybridize by forming a strong bond between the two molecules, it can be reasonably assumed that the probe and sample are essentially identical. The probe's detectable label provides a means for determining in a known manner whether hybridization has occurred. Such a probe analysis provides a rapid method for identifying formicidal δ -endotoxin genes of the subject invention.

The nucleotide segments which are used as probes according to the invention can be synthesized by use of DNA synthesizers using standard procedures. In the use of the nucleotide segments as probes, the particular probe is labeled with any suitable label known to those skilled in the art, including radioactive and non-radioactive labels. Typical radioactive labels include ^{32}P , ^{125}I , ^{35}S , or the like. A probe labeled with a radioactive isotope can be constructed from a nucleotide sequence complementary to the DNA sample by a conventional nick translation reaction, using a DNase and DNA polymerase. The probe and sample can then be combined in a hybridization buffer solution and held at an appropriate temperature until annealing occurs. Thereafter, the membrane is washed free of extraneous materials, leaving the sample and bound probe

molecules typically detected and quantified by autoradiography and/or liquid scintillation counting.

Non-radioactive labels include, for example, ligands such as biotin or thyroxine, as well as enzymes such as hydrolases or peroxidases, or the various chemiluminescers such as luciferin, or fluorescent compounds like fluorescein and its derivatives. The probe may also be labeled at both ends with different types of labels for ease of separation, as, for example, by using an isotopic label at the end mentioned above and a biotin label at the other end.

Duplex formation and stability depend on substantial complementarity between the two strands of a hybrid, and, as noted above, a certain degree of mismatch can be tolerated. Therefore, the probes of the subject invention include mutations (both single and multiple), deletions, insertions of the described sequences, and combinations thereof, wherein said mutations, insertions and deletions permit formation of stable hybrids with the target polynucleotide of interest. Mutations, insertions, and deletions can be produced in a given polynucleotide sequence in many ways, by methods currently known to an ordinarily skilled artisan, and perhaps by other methods which may become known in the future.

The potential variations in the probes listed is due, in part, to the redundancy of the genetic code. Because of the redundancy of the genetic code, *i.e.*, more than one coding nucleotide triplet (codon) can be used for most of the amino acids used to make proteins. Therefore different nucleotide sequences can code for a particular amino acid. Thus, the amino acid sequences of the *B. thuringiensis* δ -endotoxins and peptides can be prepared by equivalent nucleotide sequences encoding the same amino acid sequence of the protein or peptide. Accordingly, the subject invention includes such equivalent nucleotide sequences. Also, inverse or complement sequences are an aspect of the subject invention and can be readily used by a person skilled in this art. In addition it has been shown that proteins of identified structure and function may be constructed by changing the amino acid sequence if such changes do not alter the protein secondary structure (Kaiser and Kezdy, 1984). Thus, the subject invention includes mutants of the amino acid sequence depicted herein which do not alter the protein secondary structure, or if the structure is